

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/38, 7/01, 15/63, A61K 48/00, C07K 14/035, A61K 38/16, 39/42	A1	(11) International Publication Number: WO 95/15384 (43) International Publication Date: 8 June 1995 (08.06.95)
(21) International Application Number: PCT/CA94/00657 (22) International Filing Date: 29 November 1994 (29.11.94) (30) Priority Data: 08/159,890 30 November 1993 (30.11.93) US (71)(72) Applicants and Inventors: JOHNSON, David, C. [CA/CA]; 107 Traymore, Hamilton, Ontario L8S 1R8 (CA). YORK, Ian, A. [CA/CA]; Apartment 1207, 644 Main Street West, Hamilton, Ontario L8S 1A1 (CA). (74) Agents: MORROW, Joy, D. et al.; Smart & Biggar, Station D, P.O. Box 2999, 900-55 Melcalfe Street, Ottawa, Ontario K1P 5Y6 (CA).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: VECTOR, VIRAL PROTEIN, NUCLEOTIDE SEQUENCE CODING THEREFOR AND METHOD FOR INHIBITING IMMUNE RECOGNITION		
(57) Abstract <p>This invention relates to the use of Herpes Simplex Virus (HSV) immediate early protein ICP47, nucleic acid sequences coding for ICP47, and homologous proteins and nucleic acid sequences, to inhibit presentation of viral and cellular antigens associated with major histocompatibility class I (MHC class I) proteins to CD8+ T lymphocytes; this inhibition effectively increases infective persistence, which can, for example, improve the utility of viral gene therapy vectors. This invention also pertains to a method for the treatment of herpesvirus infections, wherein expression and/or activity of the ICP47 protein or its homologue is inhibited in order to increase immune recognition of herpesvirus-infected cells and other cells. This invention also pertains to a method for identifying drugs that interfere with the expression or function of ICP47 and its homologues, and which are useful in treating herpesvirus infections, and also pertains to the drugs so identified. Furthermore, this invention pertains to methods for the treatment and prevention of autoimmune diseases, tissue and organ transplant rejection, diabetes, multiple sclerosis, arthritis, and tissue damage accompanying ocular herpesvirus infections, wherein ICP47 or its homologue, or nucleic acids encoding such proteins, are introduced into the cells of a patient. In addition, this invention pertains to vector elements, vectors, polypeptides and polypeptide fragments that can be utilized for the foregoing purposes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo			SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Liechtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Viet Nam
GA	Gabon				

VECTOR, VIRAL PROTEIN, NUCLEOTIDE SEQUENCE CODING THEREFOR AND METHOD
FOR INHIBITING IMMUNE RECOGNITION

5

SUMMARY OF THE INVENTION

10 This invention relates generally to the use of isolated viral proteins and isolated viral nucleic acids to inhibit the ability of the immune system to recognize and then destroy virus-infected cells or other cells. This invention also relates generally to the inhibition of viral genes, mRNA and proteins in vivo in order to increase immune recognition of infected cells and other cells. More specifically, the invention relates to the use of Herpes Simplex Virus (HSV) immediate early protein ICP47, DNA sequences coding for ICP47, and homologous proteins and DNA sequences, to inhibit presentation of viral and cellular antigens associated with major histocompatibility class I (MHC class I) proteins to T lymphocytes. This inhibition effectively increases infective persistence, which can, for example, improve the utility of viral gene therapy vectors. This invention also more specifically pertains to methods for inhibiting expression and/or activity of the ICP47 protein, which can increase immune recognition of herpesvirus-infected cells and other cells, and which can, for example, serve as a means of treating herpesvirus infections.

30

BACKGROUND OF THE INVENTION

5 The normal mammalian immune system responds to viral infection in a variety of ways. One important response is that T lymphocytes become able to recognize and kill virus-infected cells, while leaving non-infected cells unharmed. Since viruses multiply by taking over the cell's machinery, when T lymphocytes kill the virus-infected cell they thereby limit the ability of the virus to reproduce itself.

10 The ability of T lymphocytes to kill only infected cells is mediated by the ability of the infected cells to produce certain "signals". These "signals", which are protein-peptide complexes called major histocompatibility (MHC) complexes, are produced by 15 mammalian cells in response to viral infection. These complexes are then transported to the surface of the infected cells, where they are "displayed" to other cells, most notably T lymphocytes. As T lymphocytes circulate in the body, they come into contact with cells 20 that have MHC complexes on their surfaces. If those MHC complexes have associated with them viral or foreign antigens in the form of small fragments of viral or foreign proteins, receptors on the surfaces of the T lymphocytes become activated, and the T lymphocytes are induced to kill those cells. But when T lymphocytes come into contact with cells that do not have the viral or foreign antigens associated with the MHC complexes on their surface, the T lymphocytes do not disturb them. 25 (Yewdell, J.W., and Bennink, J.R., Cell biology of antigen processing and presentation to major histocompatibility complex class I molecule-restricted T lymphocytes, Adv. Immunol. 52:1-123 (1992)). 30

There are two classes of MHC complexes, class I and class II. The production and display of MHC class I complexes is fairly well understood. Infected cells are able to degrade viral proteins to some extent, and short protein pieces, or peptides, are produced as a result. These peptides are transported from the nucleus or cytoplasm to the endoplasmic reticulum (ER) or to the Golgi apparatus; the ER and Golgi apparatus are convoluted, membranous intracellular organs involved in the post-translational processing of proteins, and in their transport to the cell surface. Once inside the ER or Golgi apparatus, the peptides bind to the MHC class I protein α -chains and β -2-microglobulin, to form a trimolecular complex (Townsend, A., Öhlén, C., Bastin, J., Ljunggren, H.G., Foster, L., and Kärre, K. Association of class I major histocompatibility heavy and light chains induced by viral peptides, Nature 340:443-448 (1989)). This complex is then transported to the cell surface, where it can be recognized by T lymphocyte receptors. Receptors on the surface of a particular type of T lymphocytes, known as virus-specific CD8+ T lymphocytes, specifically recognize the MHC class I complexes that are formed by the combination of MHC class I proteins and peptides derived from a particular virus, and induce the CD8+ lymphocytes to kill the cells that bear those complexes.

The presentation of MHC class I complexes and their recognition by CD8+ T lymphocytes has been also implicated in a variety of human and animal afflictions other than viral infection. Perhaps the first role identified for MHC class I complexes was their role in tissue transplant rejection, which is why they are called "Major Histocompatibility Complexes" (MHC). MHC Class I

complexes appear to be of particular importance in skin graft rejection. (Zijlstra, M., Auchincloss, H., Loring, J., Chase, C., Russell, P., and Jaenisch, R., Skin graft rejection by β_2 -microglobulin-deficient mice, J. Exp. Med. 175:885-893 (1992)). In addition, a large number of autoimmune diseases are believed to be the result of CD8+ T lymphocytes attacking cells displaying MHC class I complexes. For example, there is evidence that attack by CD8+ T lymphocytes plays a role in multiple sclerosis (see Steinman, L., Autoimmune disease Sci. Amer. 269(3):106-114), diabetes (Oldstone, M.B.A., Nerenberg, M., Southern, P., Price, J., and Lewicki, H., Virus infection triggers insulin-dependent diabetes mellitus in a transgenic model: role of anti-self (virus) immune response, Cell 65:319-331 (1991)), and arthritis (Braun, W.E., HIA molecules in autoimmune diseases, Clin. Biochem. 25(3):187-191 (1992); Scarpa, R., Del Puente, A., di Girolamo, C., della Valle, G., Lubrano, E., and Oriente, P., Interplay between environmental factors, articular involvement, and HLA-B7 in patients with psoriatic arthritis, Annals of Rheumatic Dis. 51:78-79 (1992)).

Although viral infection usually results in the display and recognition of MHC complexes, there are a number of animal viruses that are able to persist in the body, despite these mechanisms in the immune system that usually detect and destroy infected cells. Some such persistent viruses produce an extended or even constant infection, while others are able to become dormant or latent for long periods and then reappear to reinfect the individual. It is now recognized that some of these viruses evade detection by producing proteins that interfere with or block the cell's ability to make or

display MHC class I complexes (Gooding, L.R., Virus proteins that counteract host defenses, Cell 71:5-7 (1992)).

5 Different persistent viruses appear to interfere with different stages in the production and display of MHC complexes. For example, the E1a gene of adenovirus type 12 produces a protein that blocks transcription of the MHC class I genes, thus preventing the production of the MHC class I proteins themselves
10 (Schrier, P.I., Bernards, R., Vaessen, R.T.M.J., Houweling, A., and van der Eb, A.J., Expression of class I major histocompatibility antigens switched off by highly oncogenic adenovirus 12 in transformed rat cells, Nature 305:771-775 (1983)). The E3 gene of human
15 adenovirus types 2 and 5 produces a 19 thousand dalton (KD) protein that binds to the MHC class I proteins and causes them to remain sequestered or "stuck" in the ER or Golgi apparatus (Burgert, H.-G., and Kvist, S., An adenovirus type 2 glycoprotein blocks cell surface
20 expression of human histocompatibility class I antigens, Cell 41:987-997 (1985)). Similarly, murine cytomegalovirus produces a protein that inhibits the transport of the completed protein-peptide complexes from the Golgi apparatus to the cell surface (del Val, M.,
25 Hengel, H., Häcker, H., Hartlaub, U., Ruppert, T., Lucin, P., and Koszinowski, U.H., Cytomegalovirus prevents antigen presentation by blocking the transport of peptide-loaded major histocompatibility complex class I molecules into the media-golgi compartment, J. Exp. Med.
30 176:729-738 (1992)). Using an apparently very different mechanism, myxoma virus appears to cause the MHC class I proteins to be removed from the cell surface (Boshkov, L.K., Macen, J.L., and McFadden, G., Virus-induced loss

of class I MHC antigens from the surface of cells infected with myxoma virus and malignant rabbit fibroma virus, J. Immunol. 148:881-887 (1992)).

5 Herpes simplex virus (HSV) types 1 and 2 are persistent viruses that commonly infect humans; they cause a variety of troubling human diseases. HSV type 1 causes oral "fever blisters" (recurrent herpes labialis), and HSV type 2 causes genital herpes, which has become a major venereal disease in many parts of the world. No
10 fully satisfactory treatment for genital herpes currently exists. In addition, although it is uncommon, HSV can also cause encephalitis, a life-threatening infection of the brain. (The Merck Manual, Holvey, Ed., 1972; Whitley, Herpes Simplex Viruses, In: Virology, 2nd Ed., Raven
15 Press (1990)).

A most serious HSV-caused disorder is dendritic keratitis, an eye infection that produces a branched lesion of the cornea, which can in turn lead to permanent scarring and loss of vision. Ocular infections with HSV
20 are a major cause of blindness in North America. Immune responses play a major role in causing the tissue damage that results from recurrent ocular HSV infections, and T lymphocyte-mediated responses are a prominent cause of this damage. There is evidence that the CD8+ T cell
25 subset is very important in these destructive immune responses (Hendricks, R.L., and Tumpey, M., Contribution of virus and immune factors to herpes simplex virus type I-induced corneal pathology, Invest. Ophthalmol. Vis. Sci. 31:1929-1939 (1990)).

30 On initial infection, HSV usually produces a generalized, acute infection, which is cleared by the body's normal immune response. However, during the acute phase, some virus particles invade sensory nerve cells,

and there they are able to become latent, and survive long after the acute infection has been cleared by the immune system, even though antibodies against them are abundant in the blood. They then later become re-
5 activated and produce local infections. These are, as might be expected, fairly rapidly cleared by the already-prepared immune system. (Zweerink, H.J., and Stanton, L.W., Immune response to herpes simplex virus infections: virus-specific antibodies in sera from patients with
10 recurrent facial infections, Infect. Immun. 31:624-630 (1981)). This cycle is quite familiar to those who are prone to "fever blisters", which appear to be caused by sunlight-induced activation of latent HSV particles in the lips.

15 Like certain other persistent viruses, it appears that HSV inhibits immune recognition of infected cells by interfering with the synthesis, transport or display of MHC class I complexes. One reason that this was not immediately appreciated by immunologists studying
20 anti-HSV immunity is that in mouse models of HSV infection, the infected cells are primarily killed by HSV-specific CD8+ T lymphocytes, which specifically recognize MHC class I protein-HSV peptide complexes; this suggests that in these models, CD8+ T lymphocyte
25 recognition is not strongly inhibited. However, in humans, the HSV-infected cells are more often specifically killed by HSV-specific T lymphocytes of another class, called CD4+, which recognize complexes composed of HSV-derived peptides and MHC class II
30 proteins. (Schmid, D.S. and Rouse, B.T., The role of T cell immunity in control of herpes simplex virus, In: Herpes Simplex Virus: Pathogenesis, Immunobiology, and Control, B.T. Rouse, ed. (Berlin:Springer-Verlag) pp. 57-

74 (1992)). Furthermore, it has been found that human fibroblasts that are infected with HSV are not recognized and killed by HSV-specific CD8+ lymphocytes, but are killed by non-specific natural killer (NK) cells, which are not dependent on MHC class I complexes for recognition (Posavad, C.M. and Rosenthal, K.L., Herpes simplex virus-infected human fibroblasts are resistant to and inhibit cytotoxic T-lymphocyte activity, J. Virol. 66:6264-6272 (1992)). These findings suggest that recognition by CD8+ T lymphocytes is inhibited in human HSV infections.

Exactly what mechanism, what genes and what proteins might be involved in HSV's ability to suppress immune recognition has, until discovery of the present invention, remained unknown. HSV resistance to T lymphocyte recognition was known to occur within 2 to 3 hours of infection, *id.*, but MHC class I expression on the surface of HSV-infected cells was not observed to be markedly reduced until 14-20 hours after infection (Carter, V.C., Jennings, S.R., Rice, P.L. and Tevethia, S.S., Mapping of a herpes simplex virus type 2-encoded function that affects the susceptibility of herpes simplex virus-infected target cells to lysis by herpes simplex virus-specific cytotoxic T lymphocytes, J. Virol. 49:766-771 (1984)). Furthermore, other cell-to-cell propagated inactivation mechanisms have also been observed (York, I., and Johnson, D.C., Direct contact with herpes simplex virus-infected cells results in inhibition of lymphokine-activated killer cells due to cell to cell spread of virus, J. Infect. Dis. 168:1127-1132 (1993)).

The genome of herpes simplex virus type 1 is encoded on a linear, double-stranded DNA of about 152

kilobases. The HSV-1 genome has been completely sequenced. See: McGeoch, D., M.A. Dalrymple, A.J. Davison, A. Dolan, M.C. Frame, D. McNab, L.J. Perry, J.E. Scott and P. Taylor, The Complete DNA sequence of the Long Unique Region in the Genome of Herpes Simplex Virus Type 1, J. Gen. Virol. 69: 1531-1574 (1988). The genome codes for about 76 proteins, many of which have been named according to when in the infectious cycle they are produced. The protein sequences for all of the HSV-1 proteins are known, having been deduced from their corresponding gene sequences. Furthermore, many years of research has resulted in the identification of the function for many of these proteins. Nevertheless, there are still a number of proteins encoded by the HSV-1 genome that have no known function.

One of the proteins whose function has remained unknown is the immediate-early protein ICP47. Various researchers have given this protein other names, including IE12, Vmw12, and IE5. The gene for this protein is known as US 12, and is also known as α 47. The coding region of the US 12 gene is 264 base pairs long, which means that the ICP47 protein is 88 amino acids long. Although ICP47 is observed to migrate in gel electrophoresis as a protein of about 12,000 daltons, the molecular weight, as calculated from its amino acid sequence, is 9792 daltons (McGeoch, D.J., Dolan, A., Donald, S., and Rixon, F.J., Sequence determination and genetic content of the short unique region in the genome of herpes simplex virus type 1, J. Mol. Biol. 181:1-13 (1985)).

Various researchers have previously attempted to discern the function of ICP47, but prior to the present invention, without success. Deletion of the US

12 gene has been found to have no effect on infectivity (Mavromara-Nazos, P., Ackermann, M., and Roizman, B., Construction and properties of a viable herpes simplex virus 1 recombinant lacking coding sequences of the α 47 gene, J. Virol. 60:807-812 (1986)), and the most recent reported effort to determine the function of ICP47 concluded that the US 12 gene plays "no important role in the establishment and/or reactivation from latency" (Nishiyama, Y., Kurachi, R., Daikoku, T., and Umene, K., The US9, 10, 11, and 12 genes of herpes simplex virus type 1 are of no importance for its neurovirulence and latency in mice, Virology 194:419-423 (1993)).

Herpes Simplex Virus type 1 is but one member of an extended family of viruses. HSV type 2 is a close relative; its genome is "collinear" with that of HSV type 1, with "reasonable, but not identical, matching of base pairs". (Whitley, Supra at 1845). Other members of the human herpesvirus family include cytomegalovirus, varicella-zoster virus, herpes virus 6, herpes virus 7, and Epstein-Barr virus. There are also more than 50 herpesviruses that infect more than 30 other animal species (Id.), including some that infect humans.

Herpes Simplex Virus Type 2 has a gene that corresponds to the HSV Type 1 US 12 gene. It maps in the same genomic location, and produces a protein that migrates as a 12,300 dalton protein on gel electrophoresis, which is similar to the migration of ICP47 (Marsden, H.S., Lang, J., Davison, A.J., Hope, R.G., and MacDonald, D.M., Genomic location and lack of phosphorylation of the HSV immediate-early polypeptide IE 12, J. Gen. Virol. 62:17-27 (1982)). We have compared these gene sequences, and have determined that the Herpes Simplex Virus types 1 and 2 ICP47 proteins are 45%

identical at the amino acid level, and 60% homologous when one allows substitution of similar amino acids.

Varicella-Zoster Virus does not appear to have a gene corresponding to US 12 (Davison, A.J., and D.J. McGeoch, Evolutionary Comparisons of the S Segments in the Genomes of Herpes Simplex Virus Type 1 and Varicella-Zoster Virus, J. Gen. Virol. 67:597-611 (1986)), and the pseudorabies virus does not appear to contain a sequence corresponding to US 12 in the region encoding genes corresponding to other "unique stretch" (US) genes (Zhang, G., and D.P. Leader, The Structure of the Pseudorabies Virus Genome at the End of the Inverted Repeat Sequences Proximal to the Junction with the Short Unique Region, J. Gen. Virol. 71:2433-2441 (1990)). However, it is unclear whether the many other herpesviruses contain such genes.

ADVANTAGES AND SUMMARY OF THE INVENTION

It is an object of the present invention to provide a method for introducing into cells an isolated gene or other protein coding nucleic acid sequence, the expression of which will at least partially interfere with one or more mechanism involved in specific recognition by T lymphocytes.

It is also an object of the present invention to provide a method for introducing into a virus an isolated gene or other protein coding nucleic acid sequence, the expression of which will reduce immune responses to the virus so that immune suppression or destruction of virus infected cells is reduced or delayed.

It is also an object of the present invention

to provide a method for introducing into cells a protein that will at least partially interfere with one or more mechanism involved in specific recognition by T lymphocytes.

5 It is a further object of the present invention to provide a method of introducing into virus-infected cells a protein that will enable the virus to persist by at least partially avoiding recognition by T lymphocytes.

10 It is another object of the present invention to provide a new element for a gene therapy vector, and to provide an improved gene therapy vector.

15 It is also an object of the present invention to provide a method for the treatment of herpesvirus infections and to provide a method for the elimination of latent herpesviruses.

It is also an object of the present invention to provide a method for identifying drugs useful in the treatment of herpesvirus infections, and to provide drugs identified thereby.

20 An additional object of the present invention is to provide a method to suppress T lymphocyte-mediated organ or tissue transplant rejection.

25 Yet another object of the present invention is to provide a method for the treatment of T lymphocyte mediated autoimmune diseases.

Still another object of the present invention is to provide a method for the treatment of diabetes.

30 Another object of the present invention is to provide a method for the treatment of multiple sclerosis.

Yet another object of the present invention is to provide a method for the treatment of arthritis.

Another object of the invention is to provide a method for the prevention of tissue damage that occurs as

a result of immune responses to ocular herpes infections.

According to an embodiment of the invention, a method for improving the infective persistence of a virus is described. This method comprises introducing into the viral genome an isolated nucleotide sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to improve said infective persistence.

According to another embodiment of the invention, a vector element able to suppress cell recognition by cytotoxic T lymphocytes by vector-infected cells is realized that comprises an isolated nucleotide sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to suppress said recognition.

According to yet another embodiment of the invention, a vector element able to suppress cell recognition by cytotoxic T lymphocytes by vector-infected cells is realized, which comprises an isolated nucleotide sequence selected from the group of the US 12 gene from HSV Type 1, the HSV Type 2 gene encoding the IE 12 protein, nucleic acid sequences that are more than 40% homologous to the Herpes Simplex Virus Type 1 US 12 gene, and fragments of any of the foregoing that are able to suppress said recognition.

According to yet another embodiment of the invention, a vector element is realized that comprises the 714bp NruI - XhoI fragment of pRHP6, including part of the first exon, the intron, and the entire coding

sequences of ICP47.

According to another embodiment of the invention, an adenovirus vector is realized, comprising an adenovirus having the identifying characteristics of AdICP47-1.

According to yet another embodiment of the invention, a method for inhibiting cell recognition by cytotoxic T lymphocytes is described, comprising introducing into cells an isolated nucleotide sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to inhibit said recognition.

According to another embodiment of the invention, a method for inhibiting cell recognition by cytotoxic T lymphocytes is described which comprises introducing into infected cells an isolated protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to the ICP47 protein of HSV-1, and fragments of any of the foregoing that are able to inhibit said recognition.

According to yet another embodiment of the invention, a method for the treatment of herpesvirus infections is provided, which comprises the introduction into infected cells of a nucleotide sequence that is complementary to the mRNA sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and biologically active fragments of any of the foregoing, wherein the complimentary portion of said nucleotide sequence is of sufficient length to inhibit the translation of said mRNA and thereby inhibit

the production of said protein.

According to still another embodiment of the invention, a method for the treatment of herpesvirus infections is provided, which comprises the introduction
5 into infected cells of an antibody specific for a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and antigenic fragments of any of the foregoing.

10 According to yet another embodiment of the present invention, a method for identifying drugs useful in treating herpesvirus infections is provided, which comprises establishing a model cell system that expresses a protein that is selected from the group of ICP47 of HSV
15 type 1, IE 12 of HSV type 2, proteins more than 40% homologous to ICP47, and fragments of any of the foregoing that exhibit the functional characteristics of ICP47; adding amounts of candidate compounds to samples of said model cells; and testing said samples for a trait
20 different from that observed in samples to which no such compound has been added, said trait being selected from the group of suppressed synthesis of the ICP47 homologue, decreased MHC class I protein processing, and increased CTL lysis.

25 According to a further embodiment of the invention, a method is provided for the prevention and treatment of autoimmune diseases, which comprises introducing into a patient's cells a biomolecule selected
30 from the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the

protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically effective fragments of any of the foregoing.

5 Another embodiment of the invention provides a method for the prevention and treatment of tissue and organ transplant rejection, comprising introducing into the cells of said tissue or organ a biomolecule selected from the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide
10 sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically effective fragments of any
15 of the foregoing.

Still another embodiment of the present invention is a method for the prevention and treatment of diabetes, which comprises introducing into the cells of a patient a biomolecule selected from the group of an
20 isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the protein IE 12, proteins that are more
25 than 40% homologous to ICP47, and therapeutically effective fragments of any of the foregoing.

A further embodiment of the present invention is a method for the prevention and treatment of multiple sclerosis, comprising introducing into the cells of a
30 patient a biomolecule selected from the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins

that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically effective fragments of any of the foregoing.

5 An additional embodiment of the invention is a method for the prevention and treatment of arthritis comprising introducing into the cells of a patient a biomolecule selected from the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an
10 isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically
15 effective fragments of any of the foregoing.

Another embodiment of the present invention is a method for reducing immune reactions in ocular herpesvirus infections, comprising introducing into the ocular tissues of a patient a biomolecule selected from
20 the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the
25 protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically effective fragments of any of the foregoing.

BRIEF DESCRIPTION OF THE DRAWINGS

30

Figure 1(A) is a diagram of the recombinant HSV-1 virus denoted F-US5MHC;

Figure 1(B) is an autoradiogram of

electrophoretically separated ^{35}S cysteine-labeled proteins that were expressed by uninfected Daudi cells, Daudi cells infected with recombinant F-US5MHC, and Daudi cells infected with wild-type strain HSV-1(F), and which
5 were purified by immune precipitation with rabbit anti-peptide 8 (α -p8) antiserum and rabbit anti- β 2-m antiserum (α - β 2-m);

Figure 2 shows results of HSV-1-specific cytotoxic T lymphocyte lysis assays on (A) mouse
10 fibrosarcoma (MC57) cells, (B) murine SVBALB cells, or (C) normal human fibroblasts (gwfb), each having first been infected with wild type HSV-1 (F), control virus F-US5 Δ , or F-US5MHC;

Figure 3 shows autoradiograms of
15 electrophoretically separated products from pulse-chase experiments where (A) MHC class 1 α chain proteins, (B) HSV-1 or HSV-2 glycoprotein D (gD), and (C) the transferrin receptor were immunoprecipitated using
monoclonal antibodies, and samples either were or were
20 not treated with endoglycosidase H digestion prior to electrophoresis;

Figure 4 shows electrophoretic and quantitative results of pulse chase experiments in which MHC class I
products of uninfected cells and cells infected with an
25 HSV-1 mutant lacking the virion host shut-off gene were detected with two antibodies, W6/32, which recognizes only properly folded MHC class I proteins, and HC10, which recognizes both properly folded and misfolded ones;

Figure 5 shows the autoradiograms obtained
30 after electrophoresis of immunoprecipitated, pulsed and chased MHC class I α chain proteins, both with and without prior endoglycosidase H digestion, from cells that were infected with either HSV-2 strain 333 or a

mutant lacking the virion-host shut-off function, or were alternatively infected with those same viruses after they were transcriptionally inactivated by irradiation with ultraviolet light;

5 Figure 6 shows the autoradiograms obtained after electrophoresis of immunoprecipitated, pulsed and chased MHC class I α chain proteins, both with and without endoglycosidase H digestion, from cells infected with wild-type (KOS) HSV-1, and with HSV-1 mutants each
10 defective in their ability to express a single gene, either the virion host shut-off gene (vhs⁻), or one of the genes for the immediate early proteins ICP4, ICP6, ICP0, ICP22, ICP27, and ICP47;

15 Figure 7(A) is a diagram of the recombinant adenovirus vector designated AdICP47-1;

 Figure 7(B) shows the electrophoresis pattern obtained when an ICP47-specific antibody was used to precipitate radiolabeled proteins produced by cells
20 infected with wild-type HSV-1 or with adenovirus vector AdICP47-1;

 Figure 7(C) shows the autoradiograms obtained after electrophoresis of immunoprecipitated, pulsed and chased MHC class I α chain proteins, both with and without endoglycosidase H digestion, from human
25 fibroblast cells infected with wild-type HSV-1(KOS), AdICP47-1, or AddlE1 (which does not express ICP47);

 Figure 8 shows the results of human cytomegalovirus-specific cytotoxic T lymphocyte lysis assays on human MR fibroblast cells that were uninfected,
30 infected with human cytomegalovirus (CMV), or were infected with AdICP47-1 or AddlE1 followed by infection by CMV, and a similar assay on human allogeneic DG fibroblast cells that were subsequently infected by CMV.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5

It is important to an understanding of the present invention to note that all technical and scientific terms used anywhere herein, unless otherwise defined, are intended to have the same meaning as commonly understood by one of ordinary skill in the art; that techniques employed herein are also those that are known to one of ordinary skill in the art, unless stated otherwise; and that publications mentioned herein are incorporated by reference.

15

It is also important to note that reference to particular DNA fragments, genes, cDNAs, mRNAs, complementary strands, protein expression products and the like, or to some subset of such related materials (e.g., reference to DNA, where other related materials are not specifically listed) is not intended to be limiting, but should be read to include all such related materials that one of ordinary skill in the art would recognize as being of interest or value in the particular context in which that discussion is presented. It is often possible to produce or procure a biomolecule that is structurally related to or derived from a stated material, and to use that biomolecule in a different but known procedure to achieve the same goals as those to which a the use of a suggested method, material or composition is directed. For example, it is often possible to use RNA instead of DNA to carry genetic information. It is also possible to use certain nucleic acid analogues in such applications. All such

20

25

30

substitutions and modifications are included within the scope of the present invention.

It should also be noted that references to antibodies include both polyclonal and monoclonal antibodies, and also include sequences of nucleic acid that bind specifically to particular proteins of interest, which nucleic acids are referred to as "nucleic acid antibodies" (Gold, L., Nucleic Acid Ligands, PCT Application No. WO 91/19813, published Dec. 26, 1991).

Reference to a degree of homology between nucleic acids or proteins means the percentage of nucleic acid bases or amino acids that are located identically in the sequences being compared, as is commonly understood by those of ordinary skill in the art, unless it is specified that similar amino acids should be allowed. Where specified, allowing similar amino acids to be substituted means that hydrophobic amino acids may be substituted for one another, as may cationic amino acids be substituted for one another, etc., etc. Furthermore, in either case, although degrees of homology may be stated specifically, e.g., 40%, they are meant to include further levels of homology, e.g., 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% and 100%. Any mention of nucleic acid sequences encoding ICP47 or homologous proteins is intended to include all possible sequences of nucleic acids that might encode such proteins, and is not intended to be limited to the sequences derived from biological sources. Where no specific degree of homology is specified, the term "homologous" means having at least a 25% degree of homology.

In addition, when homologues of ICP47 are mentioned, it is not intended that this be limited to those homologues that occur in nature; as is known to

those skilled in the art, by using modern molecular biological techniques of site-directed mutagenesis, vector expression and the like, it is possible to produce new polypeptides, and the nucleic acid sequences that encode them, that are homologous to ICP47 and its coding sequence to various levels and in tremendously varied specific ways. It is intended that such constructs, as well as the homologues that occur in nature, be included within the scope of this invention.

It should furthermore be understood that the scope of the present invention is not limited to full-length sequences of each of the nucleic acid and amino acid sequences described. As is well understood by those skilled in the art, it is often possible to prepare subfragments of those sequences, and for those subfragments, even small ones, to retain some or all of the biological activity of the full-length sequences. Such subfragments are included within the scope of this invention.

The terms "virus" and "vector" as used herein are not intended to be mutually exclusive; to the contrary, they overlap considerably. A vector often is properly termed a virus, as that term is commonly understood. A vector often is simply a virus that has had a genetic element added, and both the term virus and the term vector would properly apply. However, a vector may have a form other than that of a virus. In addition, it is not intended that the term virus only mean replicating virus particles; the term is intended, for example, to include non-replicating virus particles, portions of viruses, and bacterial plasmids.

The term "heavy chain" is equivalent to the term "MHC class I α -chains"; H-2K^b is a specific heavy

chain derived from a particular strain of mouse.

Experimental Procedures

5 Cells and Viruses

Many of the cell types used were of a common type, and are commercially available. Some cells used were obtained commercially, and others were obtained from various laboratories. The types of cells used, the non-commercial sources from which they were obtained (if any), and some key references in which they are described are listed below:

- Vero (African Green Monkey kidney) cells;
- R-970-5 human osteosarcoma cells (Rhim, J.S., Cho, H.Y., and Huebner, R.J., Non-producer human cells induced by murine sarcoma cells, Int. J. Cancer 15:23-29 (1975)), were obtained from K. Huebner and C. Croce of the Wistar Institute, Philadelphia, Pa.;
- MC57 cells, which are mouse fibrosarcoma cells of the H-2^b haplotype (Zinkernagel, R.M., Adler, B., and Holland, J.J. Cell-mediated immunity to vesicular stomatitis virus infections in mice, Exp. Cell Biol. 46:53-70 (1978)) were obtained from M. Buchmeier of the Scripps Institute, La Jolla, CA;
- B6/WT-3 cells, which are mouse cells of the H-2^b haplotype (Pretell, J., Greenfield, R.S., and Tevethia, S.S., Biology of simian virus 40 (SV40) transplantation antigen (T Ag). V. In vitro demonstration of SV40 T Ag in SV40 infected nonpermissive mouse cells by the lymphocyte mediated cytotoxicity assay, Virology 97:32-41 (1979)) were obtained from S. Tevethia at the University of Pennsylvania, Hershey, Penn.;

- 5 - SVBALB cells, which are mouse cells of the H-2K^d haplotype (Gooding, L.R., Specificities of killing by T lymphocytes generated against syngeneic SV40 transformants: studies employing recombinants within the H-2 complex, J. Immunol. 122:1002-1008 (1979)), were obtained from K. Rosenthal at McMaster University, Hamilton, Ontario, Canada;
- 10 - Daudi cells, which are EBV-transformed human lymphoblastoid cells in which the β_2m genes are not expressed, (Klein, E., Klein, G., Nadkarni, J.S., Nadkarni, J.J., Wigzell, H., and Clifford, P., Surface IgM-kappa specificity on a Burkitt lymphoma cell in vivo and in derived cell lines, Cancer Res. 28:1300-1310 (1968));
- 15 - 293 cells, (Graham, F.L., Smiley, J., Russell, W.C., and Nairn, R., Characteristics of a human cell line transformed by DNA from human adenovirus 5 DNA. J. Gen. Virol. 36:59-74 (1977));
- 20 - Normal human fibroblasts denoted gwfb, derived from a skin biopsy (used between passages 10 - 20), EBV-transformed lymphoblastoid cell lines, obtained from K. Rosenthal of McMaster University, Hamilton, Ontario, Canada;
- 25 - The human CD8⁺ CTL clone, MR-16E6, which is specific for human cytomegalovirus (HCMV) phosphoprotein 65, was isolated and propagated as previously described (Riddell, S.R. and Greenberg, P.D., The use of anti-CD3 monoclonal antibodies to clone and expand human antigen-specific T cells, J. Immunol. Methods 128:189-201 (1990); Riddell, S.R., Watanabe, K.S., Goodrich, J.M., Li, C.R., Agha, M.E., and Greenberg, P.D., Restoration of viral immunity in immunodeficient humans by the adoptive transfer of T
- 30

cell clones, Science 257: 238-257 (1992)).

Each of the forgoing cell strains were passaged in alpha-minimal essential media (α -MEM) containing 5 to 10% fetal bovine serum (FBS), unless otherwise indicated.

Many of the viruses used were also of a common type, and are commercially available. Some viruses used were obtained commercially, and others were obtained from various laboratories. The types of viruses used, the non-commercial sources from which they were obtained (if any), and some key references in which they are described are listed below:

- HSV-1 strain F (Ejercito, P.M., Kieff, E.D., and Roizman, B., Characterization of herpes simplex virus strains differing in their effect on social behaviour of infected cells, J. Gen.Virol. 2:357-64 (1968)) was obtained from P.G. Spear at Northwestern University;
- HSV-1 strain KOS (Smith, K.O., Relationships between the envelope and the infectivity of herpes simplex virus, Proc. Soc. Exp. Biol.Med. 115:814-16 (1964)) was also obtained from P.G. Spear at Northwestern University;
- HSV-2 strain 333 (Kit, S., Kit, M., Qavi, H., Trkula, D., and Otsuka, H., Nucleotide sequence of the herpes simplex virus type 2 (HSV-2) thymidine kinase gene and predicted amino acid sequence of the thymidine kinase polypeptide and its comparison with the HSV-1 thymidine kinase gene, Biochim. Biophys. Acta 741:158-170 (1983)) was also obtained from P.G. Spear at Northwestern University;
- The HSV-1 deletion mutant VhsB, lacking the vhs gene UL41 (Smibert, C.A., and Smiley, J.R., Differential regulation of endogenous and transduced β -globin

- genes during infection of erythroid cells with a herpes simplex type 1 recombinant, J. Virol. 64:3882-94 (1990)) was obtained from J. Smiley at McMaster University, Hamilton, Ontario, Canada;
- 5 - The HSV-2 mutant lacking the vhs gene (Smibert and Smiley, unpublished) was also obtained from J. Smiley at McMaster University, Hamilton, Ontario, Canada;
- 10 - The ICP0 deletion mutant, *dlx3.1*, (Sacks, W.R., and Schaffer, P.A., Deletion mutants in the gene encoding the herpes simplex virus type 1 immediate-early protein ICP0 exhibit impaired growth in cell culture, J. Virol. 61:829-839 (1987)) was obtained from P. Schaffer at the Dana-Farber Institute, Boston, MA;
- 15 - The ICP22 deletion mutant, R325-GT⁺, (Sears, A.E., I.W. Halliburton, B. Meignier, S. Silver, and B. Roizman, Herpes simplex virus 1 mutant deleted in the α 22 gene: growth and gene expression in permissive and restrictive cells and establishment of latency in mice, J. Virol. 55:338-346 (1985)) was obtained from B. Roizman at the University of Chicago;
- 20 - The ICP6 deletion mutant, ICP6 Δ , (Goldstein, D.J., and Weller, S.K., An ICP6::*lacZ* insertional mutagen is used to demonstrate that the UL52 gene of herpes simplex virus type 1 is required for virus growth and DNA synthesis, J. Virol. 62:2970-2977 (1988)) was supplied by S. Weller of the University of Connecticut, Farmington, CT;
- 25 - The ICP47 mutant, N38, (Umene, K., Conversion of a fraction of the unique sequences to part of the inverted repeats in the S component of the herpes simplex virus type 1 genome, J. Gen. Virol. 67:1035-
- 30

1048 (1986)) was obtained from K. Umene at the Kyushu University, Fukuoka, Japan;

- The HSV-1 ICP4- mutant, d120, was propagated on complementing E5 cells (DeLuca, N.A., McCarthy, A.M., and Schaffer, P.A., Isolation and characterization of deletion mutants of herpes simplex virus type 1 in the gene encoding immediate-early regulatory protein ICP4, J. Virol. 56:558-570 (1985)), and was obtained from P. Schaffer at the

Dana-Farber Institute, Boston, MA;

- The HSV-1 ICP27 deletion mutant, 5dl1.2, was propagated on complementing 3-3 cells (McCarthy, A.M., McMahan, L., and Schaffer, P.A., Herpes Simplex virus type 1 ICP47 deletion mutants exhibit altered patterns of transcription and are DNA deficient, J. Virol. 63:18-27 (1989)) was also obtained from P. Schaffer at the Dana-Farber Institute, Boston, MA;

- The HSV-1 gD⁻ mutant F-US6kan was grown on complementing VD60 cells (Ligas, M.W., and Johnson, D.C. A herpes simplex virus mutant in which glycoprotein D sequences are replaced by β -galactosidase sequences binds to but is unable to penetrate into cells, J. Virol. 62:1486-94 (1988)).

Unless otherwise specified, and all the foregoing viruses were propagated and titered on Vero cells.

Plasmids, Viral DNA and Vectors

Many of the plasmids used were of a common type, and are commercially available. Some plasmids used were obtained commercially, and others were obtained from various laboratories. These plasmids, the non-commercial

sources from which they were obtained (if any), and some key references in which they are described are listed below:

- 5 - Plasmid pTK173, containing the HSV-1 thymidine kinase gene (Smiley, J.R., Swan, H., Pater, M.M., Pater, A., and Halpern, M.E., Positive control of the herpes simplex virus thymidine kinase gene requires upstream DNA sequences, J. Virol. 47:301-310 (1983)) was obtained from J. Smiley at McMaster University, Hamilton, Ontario, Canada;
- 10 - Plasmid pD6p, containing a lacZ gene cassette under control of the HSV-1 ICP6 promoter (Goldstein, D.J., and Weller, S.K., An ICP6::lacZ insertional mutagen is used to demonstrate that the UL52 gene of herpes simplex virus type 1 is required for virus growth and DNA synthesis, J. Virol. 62:2970-2977 (1988)) was obtained from S. Weller of the University of Connecticut, Farmington, CT;
- 15 - Plasmid pCKb, containing a EcoRI fragment including the murine H-2K^b gene inserted into plasmid pUC19 (Schönrich, G., Kalinke, U., Momburg, F., Malissen, M., Schmitt-Verhulst, A.M., Mallissen, B., Hammerling, G.J., and Arnold, B., Down-regulation of T cell receptors on self-reactive T cells as a novel mechanism for extrathymic tolerance induction, Cell 65:293-304 (1991)) was obtained from W. Jefferies at the University of British Columbia, Vancouver, British Columbia;
- 20 - Plasmid pVc β 2, containing the "a" allele of murine β 2-microglobulin gene under the control of the SV40 promoter (Daniel, F., Morello, D., Le Bail, O., Chambon, P., Cayre, Y., and Kourilsky, P., Structure and expression of the mouse β 2-microglobulin gene
- 25
- 30

- isolated from somatic and non-expressing
teratocarcinoma cells, EMBO J. 2:1061-1065 (1983)),
was also obtained from W. Jefferies at the
University of British Columbia;
- 5 - Plasmid pRHP6, containing ICP4 and ICP47 sequences
from HSV-1(KOS) (Perrson, R.H., Bacchetti, S., and
Smiley, J.R., Cells that constitutively express the
herpes simplex virus immediate-early protein ICP4
allow efficient activation of viral delayed-early
genes in trans, J. Virol. 54:414-421 (1985)), was
10 obtained from S. Bacchetti at McMaster University,
Hamilton, Ontario, Canada;
- Plasmid pS456, which contains a BamHI-MscI fragment
derived from plasmid pSS17 (Johnson, D.C., Frame,
15 M.C., Ligas, M.W., Cross, A.M., and Stowe, N.D.,
Herpes simplex virus immunoglobulin G Fc receptor
activity depends on a complex of two viral
glycoproteins, gE and gI, J. Virol. 61:2208-2216
(1988));
- 20 - HSV-1 gD- mutant F-US6KAN (Smiley, J.R., Fong, B.,
and Leung, W.-C., Construction of a double jointed
herpes simplex virus DNA molecule: inverted repeats
are required for segment inversion and direct
repeats promote deletions, Virology 113:345-362
25 (1981)).

Still other plasmids, viral DNA and vectors
used in the experiments described herein were produced by
recombinant DNA techniques that are well known in the art
(Sambrook, J., Fritsch, E.F., and Maniatis, T., Molecular
30 Cloning, a Laboratory Manual, 2nd Ed., Cold Spring Harbor
Press, 1989), and were prepared as follows:

To construct the recombinant HSV-1 expressing
murine MHC class I proteins, the 2.2 kb BamHI - NruI

fragment from pVc β 2 containing the β 2-microglobulin gene was subcloned adjacent to the thymidine kinase promoter from pTK173 and the 1.2kb SalI - PvuII fragment of pCKb containing the murine H-2K^b gene was placed under the control of the ICP6 promoter from pD6p . These β 2-microglobulin/TK and H-2K^b/ICP6 genes were then inserted into a unique NruI site in the US5 gene of pS456, producing pS5MHC. pS5gal was produced by insertion of the 4.7 kb BamHI fragment containing the ICP6::lacZ cassette from pD6p into the NruI site in pS456.

Infectious viral DNA was prepared from Vero cells infected with the HSV-1 gD- mutant F-US6KAN. Vero cells were co-transfected with F-US6KAN DNA and either plasmid pS5MHC or pS5gal, producing the viral recombinants F-US5MHC or F-US5 β , respectively, as previously described (Johnson, D.C., and Feenstra, V., Identification of a novel herpes simplex virus type 1-induced glycoprotein which complexes with gE and binds immunoglobulin, J. Virol. 61:2208-2216 (1987)).

Recombinant viruses were repeatedly plaque purified on Vero cells, in which parental F-US6KAN cannot replicate.

To construct an adenovirus vector expressing ICP47, designated AdICP47-1, the 714bp NruI - XhoI fragment of pRHP6, including part of the first exon, the intron, and the entire coding sequences of ICP47, was inserted into the EcoRV - SalI region of pCA4 (C. Addison and F.L. Graham, unpublished), which contains the left side of the adenovirus type 5 genome with a deletion spanning the E1 region, into which is inserted the human cytomegalovirus (HCMV) immediate early promoter, a polylinker, and the SV40 polyadenylation signal, so that the ICP47 coding sequences were placed next to the HCMV promoter; this produced the plasmid p47NXE1. p47NXE1

was co-transfected with plasmid pBHG10, which contains full-length adenovirus 5 sequences but without the packaging signal at the leftward side of the Ad5 genome (A. Bett and F.L. Graham, unpublished) into 293 cells.

5 Recombinant adenoviruses, in which the ICP47/HCMV promoter cassette was inserted in the E1 region and containing a deletion in the E3 region, were plaque purified on 293 cells and viral DNA was examined by restriction enzyme analysis.

10 Similarly, the control adenovirus vector Add1E1 was constructed by rescuing the plasmid pCA4 with the plasmid pJM17 (McGregory, W.J., Bautista, D.S., and Graham, F.L., A simple technique for the rescue of early region I mutations in infectious human adenovirus type
15 5, Virology 163:614-617 (1988)). The resulting virus lacked the same E1 and E3 sequences as AdICP47-1, but did not encode ICP47 or any other foreign gene.

UV-inactivation of HSV

20 HSV-2(333) or HSV-2 (333 vhs-) virus stocks were prepared by suspending Vero cells in PBS containing 1% FBS, sonicating the cells extensively, and centrifuging the material at 1000 X g for 10 minutes to remove insoluble material. The viruses were then diluted
25 in PBS containing 1% FBS to 2 to 3 ml, placed in a 60 mm dish and subjected to UV light (3 joules/cm²/sec using a bacteriostatic fluorescent tube) for 2 minutes with constant stirring while on ice. UV-inactivated viruses were shown to be unable to express any viral proteins by
30 immunoprecipitation and Western blotting; the 333 stock retained vhs activity.

Antib dies

Antibodies used, their sources, and key references describing them are:

- 5 - A hybridoma expressing monoclonal antibodies Y3, which reacts with H-2K^b complexed with β_2 -microglobulin, (Jones, B., and Janeway, C.A., Jr., Cooperative interaction of B lymphocytes with antigen-specific helper T lymphocytes is MHC restricted, Nature 292:547-549 (1981)) were obtained from the American Type Culture Collection (ATCC), Bethesda, Md;
- 10 - A hybridoma expressing monoclonal antibody W6/32 (Parham, P., Barnstable, C.J., and Bodmer, W.F., Use of a monoclonal antibody (W6/32) in structural studies of HLA-A,B,C antigens, J. Immunol. 123:342-349 (1979)), which reacts with HLA-A, B, or C complexed with β_2 -microglobulin, was also obtained from ATCC;
- 15 - Rabbit antiserum raised against a peptide from exon 8 of H-2K^b (Smith, M.H., Parker, J.M.R., Hodges, R.S., and Barber, B.H., The preparation and characterization of anti-peptide heteroantisera recognizing subregions of the intracytoplasmic domain of class I H-2 antigens, Mol. Immunol. 23:1077-1092 (1986)), which reacts with H-2K^b either complexed with, or free of, β_2 -microglobulin, was provided by B. Barber of the University of Toronto, Toronto, Canada;
- 20 - Monoclonal antibody HC10, which reacts with unfolded human HLA-B and C α -chains, and certain HLA A types (Stam, N.J., Spits, H., and Ploegh, H.L., Monoclonal antibodies raised against denatured HLA-B locus heavy chains permit biochemical characterization of certain HLA-C locus products, J. Immunol. 137:2299-
- 25
- 30

2306 (1986)), was provided by H. Ploegh at the Massachusetts Institute of Technology (MIT) in Boston, MA;

- A rabbit antiserum raised to human β_2m , which cross-reacts with mouse β_2m , was obtained from Dakopatts of Copenhagen, Denmark;

- A rabbit polyclonal antiserum directed against peptide corresponding to the C-terminus of ICP47 (Palfreyman, J.W., MacLean, J.B., Messeder, E., and Sheppard, R.C. Successful use of oligopeptides as immunogens in the preparation of antisera to immediate-early gene products of herpes simplex virus type 1, J. Gen. Virol. 65:865-874 (1984)) was obtained from H. Marsden at the Institute for Virology, Glasgow;

- LP2, a monoclonal antibody specific for HSV gD (Minson, A.C., Hodgman, T.C., Digard, P., Hancock, D.C., Bell, S.E., and Buckmaster, E.A., An analysis of the biological properties of monoclonal antibodies against glycoprotein D of herpes simplex virus and identification of amino acid substitutions that confer resistance to neutralization, J. Gen. Virol. 67:1001-1013 (1986)) was a gift of A.C. Minson at Cambridge University, Cambridge, England;

- T56/14, a monoclonal antibody specific for the transferrin receptor, was obtained from Oncogene Science, Uniondale, NY.

Radiolabeling of cells, immunoprecipitations, and endoglycosidase H digestions

Human fibroblasts or Daudi cells were metabolically labeled with ^{35}S -methionine and ^{35}S -cysteine as previously described (Johnson, D.C., and Feenstra, V.,

Identification of a novel herpes simplex virus type 1-
induced glycoprotein which complexes with gE and binds
immunoglobulin. J. Virol. 61:2208-2216 (1987)). For
pulse-chase experiments, 100 mm plates of fibroblasts
5 were labeled for 20-30 minutes with 100 μ Ci/ml each of 35 S-
methionine and 35 S-cysteine (Dupont, Dorval, Quebec) then
cell extracts were made using 1% Nonidet P40, 0.5% sodium
deoxycholate, 50 mM Tris-HCl, pH 7.5, 100 mM NaCl
(NP40/DOC buffer) containing 2 mg/ml bovine serum albumin
10 (BSA), and 1 mM phenyl methylsulfonyl fluoride (pulse) or
cells were washed and incubated in alpha-MEM containing
1% FBS for 90 minutes then cell extracts were made
(chase).

Immunoprecipitations were carried out as
15 described previously (Id.). Cell extracts were stored
overnight at -70°C and were then clarified by
centrifugation at 87,000 X g for 1 hr, then were mixed
with ascites fluids or serum and incubated on ice for 1 -
1.5 hr. Extracts to be immunoprecipitated with
20 monoclonal antibody HC10 were first heated at 70°C for 1
hour to partially denature MHC class I molecules, and
then the extracts were cooled on ice. This treatment
increased the fraction of class I molecules precipitated
with HC10. Protein A-Sepharose was added and incubated a
25 further 1.5 - 2 hours with mixing. The protein A beads
were collected by centrifugation, washed 3 - 4 times with
NP40/DOC buffer, and proteins were eluted by adding one
volume of 2 X loading buffer (4% SDS, 20% glycerol, 4% β -
mercaptoethanol and bromophenol blue) to each volume of
30 beads and heating the beads at 100°C for 5 - 10 minutes.
The stability of the MHC class I protein complex was
determined by first labeling class I proteins in
uninfected or HSV-1 (Vhs-B)-infected fibroblasts using

³⁵S-methionine and ³⁵S-cysteine (100 uCi/ml) for 1 hour, then chasing the label for 30 minutes. Cell extracts were made using NP40/DOC buffer containing 5 mg/ml BSA and 120 TIU/ml aprotinin, and these extracts were
5 incubated for 1 hour or 18 hours on ice and then immunoprecipitated using monoclonal antibodies W6/32 or HC10.

Endoglycosidase H (endo H) digestions were performed with extracts from cells that had been labeled using a pulse-chase protocol. MHC class I, gD or
10 transferrin receptors were immunoprecipitated using the appropriate antibodies, and proteins were eluted by suspending the samples in denaturing buffer (0.5% SDS, 1% β-mercaptoethanol) and boiling them for 10 minutes. Half
15 of each eluted protein sample was treated with 1000 U endo H (New England Biolabs, Mississauga, Ontario, Canada) in reaction buffer (50 mM sodium citrate) and half was incubated in reaction buffer alone, both for 3
20 hours at 37°C. The eluted proteins were then subjected to electrophoresis through 14% polyacrylamide gels for MHC class I proteins and through 8.5% polyacrylamide gels for gD and the transferrin receptor. The gels were
25 impregnated with Enlightening (New England Nuclear, Boston, MA) and exposed to X-ray film or read using a phosphorImager (Molecular Dynamics, Sunnydale, CA).

Cytotoxic T Lymphocyte lysis assays

Cytotoxic T Lymphocyte (CTL) lysis assays involving mouse cells were performed essentially as
30 previously described (Pfizenmaier, K., Jung, H., Starzinski-Powitz, A., Rollinghoff, M., and H. Wagner, The role of T cells in anti-herpes simplex virus immunity. I. Induction of antigen-specific cytotoxic T

lymphocytes, J. Immunol. 119:939-944 (1977)). Briefly, C57BL/6 (H-2K^b) or BALB/c (H-2K^d) mice were infected with 1×10^6 PFU of virus in the hind footpad. After 5 days, the mice were anaesthetized and killed, and popliteal lymph nodes were removed and crushed through stainless steel mesh. The lymphocytes were cultured for 3 days at 37°C in RPMI 1640/10% FCS/5x10⁻⁵ mM β -mercaptoethanol (CTL medium). HSV-infected target cells ($1 - 2 \times 10^4$ cells in 200 μ l MEM) were added to each well of a 96-well plate, the cells were labeled with ⁵¹Cr (Hanke, T., Graham, F.L., Rosenthal, K.L., and Johnson, D.C., Identification of an immunodominant cytotoxic T-lymphocyte recognition site in glycoprotein B or herpes simplex virus by using recombinant adenovectors and synthetic peptides, J. Virol. 65:1177-1186 (1991)), and CTLs were added at various effector:target cell ratios to a total of 200 μ l CTL medium, and were incubated for 4 hours at 37°C.

Cytotoxic T lymphocyte lysis assays involving human cells were performed using a human CD8+ CTL clone, MR-16E6, which is specific for human cytomegalovirus (HCMV) phosphoprotein 65. Human fibroblasts were infected with a recombinant adenovirus vector, AdICP47-1 or Add1E1, for 36 hours, then were subsequently infected with HCMV for 12 hours. They were then labeled with ⁵¹Cr, mixed with the CTL clone using various effector to target (E:T) ratios, and were incubated for 5 hours at 37°C.

The results of the CTL lysis assays were determined by removing and counting 100 μ l from each well to obtain experimental release (ER) of ⁵¹Cr. Maximum release (MR) was obtained by counting aliquots after treatment with 1 M HCl. In each case samples were counted in a gamma radiation counter. Total release (TR) was calculated using the equation $TR = MR + 0.5ER$. Non-

specific release (NR) was determined using cells to which no effectors had been added. Specific release (SR) was calculated using the equation $SR = (ER - NR) / (TR - NR)$.

5

Results and Conclusions

Evidence that HSV-induced inhibition of MHC Class I complex presentation is not caused by blocking synthesis of MHC proteins, and is species-specific but not MHC class I-specific.

10

Previous observations that human CD8⁺ CTL were not able to lyse HSV-infected human fibroblasts and other normal diploid cells, eg. keratinocytes (Posavad, C.M. and Rosenthal, K.L., 1992, Supra) suggested that these cells were not recognized by CTL. To examine this further with well-characterized murine CD8⁺, HSV-specific CTL, we constructed a recombinant HSV-1, F-US5MHC, which expresses murine MHC (H-2^b) class I molecules. In this construct, the murine H-2K^b gene was placed under control of the HSV-1 ICP6 promoter, and the murine β_2 -microglobulin gene was coupled to the HSV-1 thymidine kinase (tk) promoter; both of these constructs were then inserted into the HSV-1 US5 (gJ) gene, which is not required for virus replication. The structure of this clone is diagrammed in Figure 1(A). We also constructed a control virus, F-US5 β , where the HSV-1 US5 gene was interrupted with the ICP6::lacZ cassette from pD6p (Goldstein and Weller, 1988, Supra). Expression of the H-2K^b and β_2 -microglobulin proteins was then investigated by infecting Daudi cells, which do not express β_2 -microglobulin, with F-US5MHC, with wild type HSV-1 strain F, or by leaving the cells uninfected (UN). Three hours after infection the cells were labeled with ³⁵S-methionine and ³⁵S-cysteine for 2 hours, and then cell extracts were

15

20

25

30

made, and the H-2K^b α -chain was immunoprecipitated using a rabbit anti-peptide 8 (α -p8); or, the β_2 -microglobulin protein was immunoprecipitated using a rabbit anti- β_2 -m antiserum (α - β_2 -m). The results are shown in Figure 1(B); molecular mass markers are shown on the right. Expression of both the H-2^b α -chain and murine β_2 -microglobulin was detected in F-US5MHC-infected human Daudi cells, which do not normally express β_2 -microglobulin, but not in Daudi cells infected with wild-type HSV type 1 strain F (lanes marked "F"). Other experiments confirmed that the H-2K^b α -chain was expressed in human fibroblasts and a number of other human cell types infected with F-US5MHC, and that this heavy chain protein reacted with the anti-H2K^b conformation-dependent monoclonal antibody Y3 (data not shown).

F-US5MHC should in theory render any cell susceptible to lysis by murine, H-2^b-restricted CTL. Figure 2 shows that when cytotoxic T lymphocytes (CTL) derived from C57BL/6 mice (H-2^b) infected with HSV-1(F) were used in CTL assays using effector to target ratios of 40, 20, or 10:1, mouse fibrosarcoma H-2^b (MC57) target cells infected with F-US5MHC were efficiently lysed (Figure 2(A)), as were (murine H-2^d) SVBALB cells (Figure 2(B)). However, uninfected (UI) MC57 cells were not lysed (Figure 2(A)), nor were uninfected SVBALB cells or SVBALB cells infected with wild type HSV-1(F) or infected with control virus F-US5 β . In other experiments, rat cells infected with F-US5MHC were also rendered susceptible to lysis by H-2^b-restricted CTL (data not shown). In contrast, normal human fibroblasts (gwf^b) (Figure 2(C)) and a panel of other human cells (data not shown) were not lysed by HSV-specific, H-2^b-restricted CTL after infection with F-US5MHC; nor were uninfected (UI)

human normal fibroblasts or fibroblasts infected with F-US5 β or F-US5MHC. Therefore, these human cells were not recognized by mouse cytotoxic T lymphocytes even though they expressed mouse MHC class I molecules. Together
5 these results suggest that the HSV-induced inhibition of presentation to CTL is not related to inhibition of MHC class I synthesis and may be species-specific, but is not MHC class I-specific.

10 **Evidence that MHC class I molecules in HSV-infected cells are retained within the ER/cis Golgi compartment**

To further study MHC class I molecules in HSV-infected human cells, we used a pulse-chase protocol to examine intracellular transport and processing of class I
15 α -chain molecules. Normal human fibroblasts (gwfb) were left uninfected or were infected with HSV-1(KOS) or HSV-2(333) for 3 hr, then labeled with 35 S-methionine and 35 S-cysteine for 30 minutes and lysed (pulse: P) or the label was chased for 90 minutes (chase: C) before lysis. The
20 results are shown in Figure 3. In Figure 3(A), MHC class I α chain proteins were immunoprecipitated using monoclonal antibody HC10; in Figure 3(B), HSV-1 or HSV-2 glycoprotein D (gD) were immunoprecipitated using
25 monoclonal antibody LP2; and in Figure 3(C), the transferrin receptor was immunoprecipitated using the monoclonal antibody T56/14. The proteins were eluted from protein A beads and digested with endo H (+) or mock digested (-) at 37 °C before electrophoresis and
30 autoradiography. In Figure 3(D), human fibroblasts were infected with HSV-1 (F) for 0, 2, 4, 6, or 8 hours, were then pulse labeled for 30 minutes as in (A), and the label was chased for 90 minutes. MHC class I molecules were immunoprecipitated with antibody W6/32 and samples

were either treated (+) or not treated (-) with endo H. A molecular mass marker of 45 KDa is indicated on the right.

5 The results show that Class I molecules
immunoprecipitated from infected or uninfected cells were
digested with endoglycosidase H (endo H), which removes
high-mannose but not fully processed oligosaccharides, as
a measure of glycoprotein transit through the medial and
trans Golgi compartments (Townsend et al, 1989. Supra).
10 MHC class I α -chains from uninfected cells became
resistant to endo H after a 90 minutes chase period,
while class I proteins from cells infected with HSV-1 or
HSV-2 remained sensitive to endo H (Figure 3A). The
inhibition of MHC class I transport and processing in
15 HSV-infected cells was apparently a specific effect
rather than a general one, since HSV-1 glycoprotein D
(gD) and the transferrin receptor were efficiently
processed to become endo H-resistant during the 90
minutes chase period (Figure 3B,C). When cells were
20 infected with HSV-1 and examined at various times after
infection, alterations in the processing of MHC class I
was first observed 2 hours after infection with HSV-1 and
the effect was complete by 4 hours (Figure 3D). As
expected, mouse H-2K^b class I molecules expressed in human
25 fibroblasts infected with F-US5MHC also remained in an
endo H sensitive form, yet H-2K^b and several other mouse
MHC class I proteins expressed in HSV-infected mouse
fibroblasts and other mouse cells became endo H resistant
(data not shown). These results demonstrate that MHC
30 class I complexes are retained in the endoplasmic
reticulum/cis Golgi of human fibroblasts, but not mouse
cells, soon after infection with either HSV-1 or HSV-2.

It is well established that processing of N-

linked oligosaccharides occurs in the Golgi apparatus as glycoproteins are transported from the endoplasmic reticulum (ER) (site of synthesis and initial glycosylation) to the Golgi then to the cell surface. When glycoproteins do not become processed they do not reach the cell surface. Therefore, lack of processing indicated by endo H sensitivity (endo H recognizes immature, high mannose N-linked oligosaccharides but not mature complex N-linked oligosaccharides) is indicative of lack of transport to the cell surface (for review see: Kornfeld, R., and Kornfeld, S., Assembly of asparagine-linked oligosaccharides, Ann. Rev. Biochem. 54:631-664. (1985)).

Evidence that MHC I in HSV-infected cells is unstable.

MHC class I polypeptides produced in RMA-S and T2 cells, lacking the putative peptide transporter proteins, were found to be misfolded and unstable (Townsend et al, 1989, Supra; Townsend, A., Elliott, T., Cerundolo, V., Foster, L., Barber, B., and Tse, A., Assembly of MHC class I molecules analyzed in vitro, Cell 62:285-295 (1990)). In order to examine the stability and folding of class I molecules in HSV-infected human fibroblasts, we carried out pulse-chase experiments in which we detected the MHC class I products with two antibodies, one which recognizes only properly folded MHC class I proteins, and another which recognizes both properly folded and misfolded ones.

More specifically, in these experiments human fibroblasts were left uninfected or infected for 4 hours with HSV-1 VhsB, a mutant lacking the virion host shutoff gene, then the cells were radiolabeled for 1 hour with ³⁵S-methionine and ³⁵S-cysteine and the label was chased

for 30 minutes. Cell extracts were mixed with antibodies immediately (1 hr) or were incubated for 18 hours on ice before being mixed with antibodies. MHC class I proteins were immunoprecipitated using monoclonal antibody W6/32, which recognizes only properly folded class I heavy/light chain complexes (Parham et al, 1979, Supra) or HC10 which, under the conditions used, recognizes misfolded as well as folded MHC class I α -chains (Stam et al, 1986, Supra), by first heating the cell extracts for 1 hour at 70°C to denature the protein molecules. After immunoprecipitation, samples were subjected to electrophoresis on 14% polyacrylamide gels, as shown in Figure 4(A); a molecular weight marker of 45 KD is shown at the right of the gel. A densitometric quantitation of protein bands corresponding to the class I α -chains was also performed, as shown in Figure 4(B). The densitometric values obtained with uninfected and HSV-infected cell extracts incubated for 1 hours and precipitated with HC10 were set at 100.

As is apparent from Figure 4, There was a modest inhibition of MHC class I α -chain synthesis in HSV-infected cells, even though an HSV-1 mutant unable to express the virion host shut-off function, vhs, was used (Smibert and Smiley, 1990, Supra); perhaps this was because of competition between cellular and viral transcription and translation factors. Densitometric quantitation of the protein bands immunoprecipitated by HC10 showed that there was no appreciable proteolytic degradation of the α -chain in either infected or uninfected cells during the 18 hours incubation at 4°C. However, only a fraction of the class I α -chains present in extracts from HSV-infected cells were recognized by W6/32. In the example shown, approximately 42% of the

class I molecules precipitated by HC10 (total number of molecules) were recognized by W6/32 after 1 hours and about 30% of these molecules dissociated during 18 hours at 4°C (Figure 4). In contrast, class I molecules from uninfected cells were efficiently recognized by W6/32 and were stable, as less than 5% of the molecules dissociated during the 18 hours incubation. Therefore, it appears that MHC class I complexes formed in HSV-infected cells were misfolded and considerably less stable than those formed in uninfected cells.

In these experiments, β 2-microglobulin levels were not dramatically altered by HSV infection (data not shown), and furthermore, this β 2-microglobulin was available for binding to class I heavy chains because 45% of the heavy chain could be recognized by monoclonal W6/32, which recognizes only class I complexes containing β 2-microglobulin.

It is known that folding of MHC class I proteins in the ER is dependent upon trimerization of MHC class I heavy or alpha chain, β 2-microglobulin, and small peptides derived from cellular or viral proteins (reviewed in Yewdell and Bennick (1990), Supra). Townsend et al. (1990, Supra) and numerous others have shown that MHC class I molecules fail to assemble, fold improperly and are not transported to the cell surface in mutant cells if peptides are not available in the ER. Later studies indicated that this was because these mutant cells lack TAP proteins which "pump" peptides into the ER. In these mutant cells, eg. RMA-S or .174, MHC class I proteins remain sensitive to endo H and are misfolded. Moreover, the observed misfolding and instability of the MHC class I complexes in HSV-infected fibroblasts is similar to that observed in TAP

transporter-negative cell lines, and indicates that peptides are not associated with these MHC class I complexes in HSV-infected cells.

5 It has been found here that MHC class I proteins synthesized in HSV-infected cells have the same attributes, e.g., the class I proteins remain endo H sensitive and are misfolded, as indicated by their lack of recognition by a conformationally sensitive monoclonal antibody. Since their lack of processing means that the class I proteins do not reach the cell surface in HSV-10 infected cells, one predicts that the class I proteins would be defective in presenting viral antigens to T lymphocytes.

15 **Evidence that the HSV-1 immediate-early gene product ICP47 is required for ER retention of MHC I.**

HSV expresses three classes of gene products: immediate early (IE), early (E), and late (L), where IE proteins are required for the synthesis of E and L proteins (Honess, R.W. and Roizman, B., Regulation of herpesvirus macromolecular synthesis. I. Cascade regulation of the synthesis of three groups of viral proteins, J. Virol. 14:8-19 (1974)). However, a group of viral gene products including the vhs protein (McLaughlin, J., Addison, C., Craigie, M.C., and Rixon, F.J., Noninfectious L-particles supply functions which can facilitate infection by HSV-1, Virology 190:682-688 (1992)) and the VP16 transactivator of IE proteins (Batterson, W., and B. Roizman., Characterization of the herpes simplex virion-associated factor responsible for the induction of α genes, J. Virol. 46:371-377 (1983)) are incorporated into the virus particle and delivered into host cells upon virus entry. Since MHC class I

20
25
30

proteins were retained in the ER within 2 hours following HSV-1 infection (Figure 2), it appeared that either a virion structural protein or an immediate-early gene product was be responsible for the retention of those proteins.

To determine whether a virion structural protein was involved in this effect, stocks of HSV-2 were subjected to UV-inactivation so that the virus particles retained vhs activity but were transcriptionally silent. Human fibroblasts were left uninfected, infected with HSV-2 (333), or with HSV-2 (333-vhs⁻), a mutant derived from 333 which does not express the vhs function, using 10 plaque forming units/ml (PFU/ml). Other monolayers of fibroblasts were treated with gradient purified, UV-inactivated virus particles derived from HSV-2 strain 333 or 333-vhs⁻ at levels corresponding to 200 PFU/cell, and were incubated for 2 hours at 37 °C . The cells were labeled using the pulse-chase protocol described for the experiments shown in Figure 3, except that the pulse was for 20 min; then MHC class I proteins were immunoprecipitated using antibody W6/32, and class I proteins digested (+) or not digested (-) with endo H, as also described for the experiments shown in Figure 3. The proteins were then subjected to electrophoresis on 14% polyacrylamide gels and exposed to X-ray film.

The results are shown in Figure 5. In cells treated with relatively large quantities of UV-inactivated HSV-2 particles lacking the vhs protein (333-vhs⁻), MHC class I proteins were processed in an identical fashion to that in uninfected cells; processing to endo H resistant forms after the chase period was not significantly inhibited. UV-inactivated virus particles derived from a HSV-2 strain which retained a wild type

5 vhs gene (333-vhs+) produced a marked decrease in the expression of class I proteins under these conditions, demonstrating that the virus particles retained vhs activity after UV-inactivation which had shut off host transcription.

10 Since placing relatively high concentrations of structural proteins in the presence of cells was inadequate to inhibit the processing of MHC class I proteins to their endo H resistant forms, it appears that HSV-2 structural proteins and the vhs protein, which are part of the virus particle, are not sufficient to cause ER retention of MHC class I proteins, and that transcription of HSV genes is required.

15 In order to determine whether HSV IE proteins were capable of inhibiting processing of MHC class I proteins, a panel of HSV-1 mutants unable to express the 6 IE proteins was analyzed. Human fibroblasts were left uninfected (UN) or were infected with wild-type HSV-1 (KOS), an HSV-1 lacking the virion host shutoff gene (KOS-vhs⁻); HSV-1 mutant dl120, which is unable to express ICP4; HSV-1 mutant ICP6 Δ , which is unable to express ICP6; HSV-1 mutant R325- β TK⁺, which is unable to express ICP22; HSV-1 mutant 5dl1.2, which is unable to express ICP27; or HSV-1 mutant N38, which is unable to express ICP47. Cells were infected for 3 hr, were labeled with ³⁵S-methionine and ³⁵S-cysteine for 20 min, and were then immediately lysed (P) or the label was chased for 90 minutes (C) before lysis. MHC class I α -chains were immunoprecipitated using monoclonal antibody HC10 and the proteins either were treated with endo H (+) or were not treated (-) before electrophoresis and autoradiography.

30 The results of these experiments are shown in Figure 6. In cells infected with HSV-1 dl120, a mutant

unable to express IE protein ICP4, class I proteins remained endo H sensitive. Since ICP4 is strictly required for expression of both E and L proteins (Dixon, R.A.F., and Schaeffer, P.A., Fine-structure mapping and functional analysis of temperature-sensitive mutants in the gene encoding the herpes simplex virus type 1 immediate early protein VP175, J. Virol. 36:189-203 (1980); Watson, R.J. and Clements, J.B., A herpes simplex virus type 1 function continuously required for early and late virus RNA synthesis, Nature 285:329-330 (1980)), this shows that expression of only the HSV-1 IE proteins is sufficient to inhibit class I processing and transport. Analysis of MHC class I proteins produced by mutants unable to produce IE proteins ICP6, ICP22, ICP27 and ICP0 (Figure 6) shows that these mutants were still able to inhibit class I processing, as the proteins remained endo H sensitive. This indicates that these IE proteins are not essential to the inhibition of MHC class I protein processing.

In contrast, mutant N38, which is unable to express ICP47 (Nishiyama et al. (1993), Supra), did not block class I processing; the chased samples showed little or no endo H sensitivity, indicating that processing was completed or nearly completed by that time. We note that mutant N38 lacks coding sequences for the US9, US10, and US11 genes in addition to the lack of ICP47 coding sequences; however, The US9, US10 and US11 genes, which are early and late genes, were not expressed in cells infected with mutant d120, which lacks the ICP4 gene function that is required for expression of early and late genes, and cells infected with this mutant still caused inhibition of MHC Class I processing (see above). Thus, it is unlikely that any of these genes are

involved in this effect. In contrast, these results indicate that IE protein ICP47 is involved in inhibition of MHC class I protein processing and transport to the cell surface.

5 A second HSV-1 mutant, R3631, which lacks the ICP47 and US11 genes (Mavromara-Nazos, (1986), Supra), also failed to inhibit class I processing (not shown). ICP47 is an IE gene which does not require ICP4 for its expression, and it has no known effect on virus
10 replication. Expression of the other IE genes, as well as E and L proteins, in cells infected with this mutant was normal (data not shown). This indicates that ICP47 is required for the observed inhibition of class I processing and transport, and that no other IE proteins
15 are necessary.

Since MHC class I molecules must be transported to the cell surface so that cells can be recognized by CD8+ T lymphocytes, the observed inhibition of class I processing and transport caused by ICP47 would be
20 expected to cause cells containing ICP47 to be resistant to these cytotoxic T lymphocytes.

As a final and conclusive proof that the gene for ICP47 alone is sufficient to produce the inhibition of MHC class I protein processing and transport, and that
25 it is also sufficient to produce the inhibition of CTL lysis that is observed in HSV infections, the ICP47 gene has been cloned into an adenovirus vector. Studies on this vector have demonstrated that a vector carrying the ICP47 gene is able to produce the ICP47 protein, cause
30 inhibition of MHC class I protein processing and transport, and inhibit cell lysis by cytotoxic T lymphocytes. Because these experiments are of great value in illustrating many of the uses for the ICP47 gene

and its homologues that are part of this invention, they have been omitted here, and have instead been shown and described as part of Example 1 hereinbelow.

5 **Improving the infective persistence of a mammalian virus by inserting genes for ICP47 or ICP47 homologues.**

10 The use of viruses to serve as vectors for "gene therapy" is a promising technique, which may in the future be widely practiced. Simply stated, a gene
15 encoding a protein having therapeutically desired effects is cloned into a viral expression vector, and that vector is then introduced to the target organism. The virus infects the cells, and then produces the protein sequence in vivo, where it has its desired therapeutic effect.
20 (See, e.g., Zabner, J., Couture, L.A., Gregory, R.J., Graham, S.M., Smith, A.E., and Welsh, M.J., Adenovirus-mediated gene transfer transiently corrects the chloride transport defect in nasal epithelia of patients with cystic fibrosis, Cell 75:207-216 (1993)). However, one limitation of this use of viral vectors for gene therapy is that the infected cells get recognized and killed by the T lymphocytes, which eventually results in the loss of the virus and its beneficial therapeutic effect. Indeed, the concern that immune responses directed to
25 viral proteins or to foreign gene therapy proteins might cause loss of gene therapy vectors is a major concern in the field of gene therapy (Id.).

30 The present invention offers a method by which the infective persistence of viral gene therapy vectors might be improved, thus enhancing the therapeutic value of those vectors. This method entails adding to the vector a sequence that encodes the ICP47 protein, the IE 12 protein of HSV-2, or another protein substantially

homologous to ICP47. Once these sequences are added to the vector, the expression of those sequences should effectively inhibit CD8+ T lymphocyte recognition of infected cells, and should thereby enhance the longevity and virility of the beneficial virus.

One embodiment of this aspect of the invention is shown in Example 1 hereinbelow, wherein the ICP47 gene has been isolated and inserted into an adenovirus vector. This vector is thereafter able to inhibit the processing and transport of MHC class I proteins, and to thereby inhibit cell lysis by cytotoxic T lymphocytes that are directed against a virus with which the cells were co-infected. This demonstrates that such a vector or vector element could be used to protect the cells in which they reside from CTL-mediated cell lysis, whether the desired gene therapy gene was carried on that vector or on another vector with which the recipient was to be co-infected. Since it is the ICP47 gene product that causes the inhibition, these experiments suggest that delivering the ICP47 protein or a protein substantially homologous to it into the cells would have a similar effect. Such proteins could be produced using known molecular biology methodologies (See: Sambrook et al, 1989, Supra).

Of course, the foregoing should not be viewed as limited to gene therapy vectors, as this method could be used to improve persistence in any virus in which such persistence may be desired.

The present invention also includes the vector elements that might be used to carry out this method. It is possible to use only the protein coding regions for such purpose, or to use the entire genes, including introns and other elements. The 714bp NruI - XhoI fragment of pRHP6, including part of the first exon, the

intron, and the entire coding sequences of ICP47, is one such element. Adenovirus vector AdICP47-1 is a gene therapy vector that includes this element.

5 **Inserting genes for ICP47 or ICP47 homologues into somatic genes or secondary vectors to inhibit cell recognition**

As an alternative to adding the sequences encoding ICP47 or a homologous protein to the DNA of a virus, it is also possible to introduce such a gene into the somatic DNA of infected or uninfected cells, by methods that are well known in the art (Sambrook et al. 1989, Supra). Alternatively, it is also possible to introduce such a sequence into infected or uninfected cells by use of a secondary virus, i.e., one that is not the virus that is expected to produce the immune response. For example, it may be desirable to introduce such a secondary virus carrying an ICP47 coding sequence prior to the introduction of a viral gene therapy vector. This possibility is well illustrated by Example 1 hereinbelow; the gene for ICP47 is carried on an adenovirus vector, yet is demonstrably able to inhibit cell lysis by cytotoxic T lymphocytes specific for cytomegalovirus, with which the cells were co-infected. Thus, it is clear to one of ordinary skill in the art that one vector carrying the gene for ICP47 or a homologous protein could be used to infect a patient, and that a second vector, e.g., a virus carrying a gene to be introduced by gene therapy, could be used to infect that same individual; the first vector would inhibit recognition by CTL directed against either vector.

Additionally, it may be quite useful to introduce a sequence that codes for ICP47 or a homologous

protein into the cells of individuals suffering from autoimmune diseases, either by introducing it into the somatic DNA or into a viral gene therapy vector, or alternatively, to introduce the protein ICP47 or its homologue into the patient's cells. By doing so, it should be possible to limit the display of MHC Class 1 complexes, and thereby limit autoimmune responses and symptoms of the disease. This approach should be useful in the treatment of a number of disorders believed to be in part mediated by cytotoxic T lymphocytes, such as tissue and organ transplant rejection, diabetes, multiple sclerosis and arthritis, whether or not they are wholly "autoimmune disorders" as that term is generally used. By reducing the recognition of cells of involved tissues by cytotoxic T lymphocytes, the symptoms of the disorder should be reduced. It should also be of value to use such a therapy with patients at high risk for developing such disorders, even if the damaging symptoms have not yet appeared. Furthermore, this approach should also be of value in the treatment of ocular herpesvirus infections, where significant tissue damage occurs as a result of cytotoxic T lymphocyte activity; this approach should lead to a reduction in recognition of cells by T lymphocytes, and thereby reduce tissue damage. Similarly, it should be possible to use this method to treat transplant recipients in order to limit CD8+ T lymphocyte-mediated tissue rejection.

In some of the foregoing examples, it may only be necessary to introduce the genetic or protein elements into certain cells or tissues. For example, in the case of diabetes, introducing them into only the pancreas should be sufficient, and in the case of tissue or organ transplants, to only introduce them into the tissues or

organ being transplanted. However, it may be more therapeutically effective and more simple to treat all of the patients cells.

5 Each of the above objectives should be equally attainable by adding to the cells the ICP47 protein itself, or a homologous protein, which could be easily produced by known recombinant DNA methodologies. (Sambrook et al., 1989, Supra).

10 **Treating persistent herpesvirus infections by use of antisense strands targeted to a ICP47 homologous mRNA or gene.**

15 There is now a considerable art regarding the use of so-called "anti-sense" polynucleotide sequences or analogs to prevent the expression of proteins in vivo (for review see: Neckers, L. and Whitesell, L., Antisense Technology: biological utility and practical considerations, Am. J. Physiol. 265 (Lung Cell. Mol. Physiol. 9:) L1--L12 (1993)). The basic theory is that
20 if you add to a cell a large number of strands of a nucleotide sequence that is complementary to the messenger RNA that is transcribed to produce a particularly protein, these "anti-sense" strands will hybridize to the mRNA and limit or prevent its
25 transcription. This method could be used here to limit or prevent the expression of ICP47 and homologous proteins by herpesviruses; as a result, suppression of immune recognition by CD8+ T lymphocytes would be reduced, and the organism's immune system could more
30 rapidly and effectively kill the infected cells. It is possible that this will even make it possible to eliminate the recurring symptoms of herpesvirus infection; e.g., in HSV-1 infections, it might be

possible to prevent recurring "fever blisters".

5 **Treating herpesvirus infections with antibodies against
an ICP47 homologous protein**

10 Persistent herpesvirus infections may also be
advantageously treated by introducing monoclonal or
polyclonal antibodies to the ICP47-homologous protein
that the given virus produces; this will limit or prevent
the suppression of MHC Type I complex expression, and
thus allow for more effective immune clearing. Methods
for intracellular introduction of antibodies have been
described (See, e.g. Carlson, J.R., A new use for
15 intracellular antibody expression: inactivation of human
immunodeficiency virus type 1, Proc. Natl. Acad. Sci. USA
90:7427-7428 (1993)).

20 **Treating herpesvirus infections with drugs which
interrupt or inhibit ICP47**

ICP47 appears to block availability of cellular
or viral peptides which would normally be presented on
MHC class I molecules, causing the MHC class I molecules
to become misfolded and preventing their transport to the
25 cell surface. Whatever the mechanism of ICP47 action,
however, it is reasonable to conclude that ICP47 achieves
its effect by interaction with one or more intracellular
protein or other molecule. Furthermore, it is clear that
inhibition of MHC class I protein display depends on
30 synthesis of ICP47, and it seems reasonable to conclude
that its effectiveness is to at least some extent dose-
dependent.

With these things in mind, it seems clear that

by blocking the synthesis of ICP47 or its homologues, or by interfering with the interaction between ICP47 or its homologues and cellular molecules using drugs specifically developed for this purpose, one would expect
5 that cells infected with herpesvirus would be more readily recognized by anti-herpesvirus CD8+ T lymphocytes, leading to better recognition of herpesviruses by the immune system, with the beneficial results of reduced infection, decreased latency, and
10 reduced symptoms.

The invention as described herein includes a method to screen for such drugs, as well as the drugs so identified. Quite simply, one need only create a system that produces ICP47 or a homologue of it, add amounts of
15 candidate compounds to that system, and determine whether synthesis of ICP47 or the homologue is inhibited, whether inhibition of the processing of MHC class I proteins decreases, or whether cytotoxic T lymphocyte lysis of cells increases when compared to that same system in the
20 absence of the added compound. Candidate compounds could include a wide spectrum of small molecules from which so-called "ethical pharmaceuticals" are often identified, and could also include a wide variety of other compounds, including large and small synthetic compounds, as well as
25 many naturally-occurring or man-made biomolecules, including polynucleotides and polypeptides.

One could easily use one or more of the methods described herein to accomplish this. For example, one might establish a model system that produces ICP47,
30 either because it carries the coding gene in its genome, on a vector, or in a virus with which it has become infected. One could then test for decreased ICP47 synthesis by using the ICP47 antibody detection method

used to produce the results shown in Figure 7; test for an increase in endo H resistance of the MHC class I complexes by using a pulse-chase protocol as was used to produce the results shown in figure 6; and/or test for increased CTL lysis using the assay method used to produce the results shown in Figure 2 or Figure 8.

EXAMPLE 1

Construction and characterization of a recombinant adenovirus vector carrying the coding sequence for HSV-1 ICP47

To determine whether other HSV gene products were required for the inhibition of antigen processing and whether ICP47 was sufficient for this effect, we constructed a recombinant adenovirus vector, AdICP47-1. As shown in Figure 7(A), the HSV-1 ICP47 gene was placed under the control of the HCMV immediate early promoter and was inserted into the E1 region of adenovirus type 5 to produce AdICP47-1. AdICP47-1 is replication defective because it lacks Adenovirus E1 sequences as well as E3 sequences.

Expression of ICP47 proteins

The expression of ICP47 was examined by infecting human fibroblasts with HSV-1(F) using 10 PFU/cell or AdICP47-1 (Ad47) using either 10 or 100 PFU/cell, or leaving fibroblasts uninfected (UN). Cells were labeled for 2 hr, HSV-infected cells were radiolabeled from 1 to 3, 2 to 4, or 3 to 5 hours after infection, and AdICP47-1-infected cells were radiolabeled

from 36 to 38 hours after infection using ^{35}S -methionine and ^{35}S -cysteine (50 $\mu\text{Ci}/\text{ml}$ of each). ICP47 was immunoprecipitated using an ICP47-specific antipeptide serum and subjected to electrophoresis using 16% polyacrylamide gels. As shown in Figure 7(B), proteins of the same molecular weight as ICP47 were produced by AdICP47-1 (lanes are marked "Ad47" in the figure legend), and these reacted with ICP47-specific antibodies. It therefore appears that the adenovirus vector AdICP47-1 is able to produce the ICP47 protein in vivo.

MHC class I transport and processing is inhibited

In order to determine whether the ICP47 proteins produced upon infection of cells with adenovirus vector AdICP47-1 were able to inhibit intracellular MHC class I protein processing, human fibroblasts were left uninfected or were infected with HSV-1(KOS), AdICP47-1, or Add1E1 (which lacks E1 and E3 sequences but does not express ICP47). Four hours after infection with HSV-1 or 36 hours after infection with an adenovirus, the cells were radiolabeled with ^{35}S -methionine and ^{35}S -cysteine for 30 minutes and immediately lysed (P) or the label chased for 90 minutes (C) before lysis. MHC class I α -chains were immunoprecipitated using monoclonal antibody HC10 and treated with endo H (+) or not treated (-) before electrophoresis and autoradiography. The results are shown in Figure 7(C); a molecular weight marker of 45 kDa is shown at the right.

These results show that MHC class I proteins produced in human fibroblasts infected with AdICP47-1 remained sensitive to Endo H, as with HSV-1-infected human fibroblasts. Class I molecules from cells infected with control adenovirus Add1E1, lacking E1 and E3

sequences, became largely endo H resistant. These results show that the ICP47 protein produced by the recombinant adenovirus vector AdICP47-1 is able to inhibit MHC class I protein processing in vivo.

5 Similar results were obtained with another adenovirus vector, AdICP47-3, where ICP47 coding sequences were placed under control of the SV40 early promoter and inserted into the E3 region so that the E1 sequences were intact and the adenovirus vector was
10 capable of replicating in human cells (data not shown). Therefore, other HSV gene products are not required for this effect and expression of HSV ICP47 is sufficient to prevent MHC I transport and processing.

15 **Lysis by specific cytotoxic T lymphocytes is inhibited**

 In order to determine if ICP47 produced by an adenovirus vector could inhibit recognition by cytotoxic T lymphocytes, human MR fibroblasts were left uninfected (UN) or were infected with AdICP47-1 or Add1E1 for 36
20 hours, and were then infected with human cytomegalovirus (CMV) for 12 to 16 hours. An allogeneic fibroblast cell line, DG, was also infected with CMV. The fibroblasts were then loaded with the radiolabel ⁵¹Cr and mixed with various ratios (effector:target cell ratios or E:T
25 ratios) of a human cytomegalovirus-specific cytotoxic T lymphocyte clone, MR-16E6. Release of ⁵¹Cr was determined after 5 hours, and percent specific lysis of the fibroblasts was calculated.

 The results are shown in Figure 8. Uninfected
30 fibroblasts (UN) were not lysed by the HCMV specific clone (I.E., no ⁵¹Cr was released after mixing with the cytotoxic T lymphocytes) but 40% to 50% of the ⁵¹Cr was released from HCMV-infected fibroblasts (CMV) indicating

that there was specific lysis of these cells. Lysis of
allogeneic DG fibroblasts (Allo + CMV), which do not
share the same MHC class I molecules as the cytotoxic T
lymphocytes, did not occur, indicating that the cytotoxic
5 T lymphocyte clone lysed only target cells which shared
the same MHC class I molecules. Prior infection of the
fibroblasts with AdICP47-1 caused the lysis of
fibroblasts subsequently infected with HCMV (AdICP47 +
CMV) to be reduced to background levels. By contrast,
10 prior infection of cells with AddlE1 had no effect on the
lysis of cells infected with HCMV (AdICP47 + CMV).
Therefore, expression of ICP47 blocked lysis of the cells
by a cytotoxic T lymphocyte clone.

15 It is understood that the examples and
embodiments described herein are for illustrative
purposes only, and that various modifications or changes
in light thereof will be suggested to persons skilled in
the art; these are to be included within the spirit and
20 purview of this application and scope of the appended
claims.

We claim:

1. A method for improving the infective persistence of a virus, comprising introducing into the genome of said virus an isolated nucleotide sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to improve said infective persistence.
2. A method for inhibiting cell recognition by cytotoxic T lymphocytes, comprising introducing into said cells an isolated nucleotide sequence encoding a protein selected from the group of ICP47 of HSV type 1; IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to inhibit said recognition.
3. The method according to claim 2 wherein the nucleotide sequence is introduced through the use of a viral vector.
4. The method according to claim 2 wherein the nucleotide sequence is introduced through the use of a viral vector separate from and in addition to a gene therapy vector with which the cells are also infected.
5. A vector element able to suppress recognition of vector-infected cells by cytotoxic T lymphocytes, comprising an isolated nucleotide sequence encoding

a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to suppress said recognition.

6. A vector element according to claim 5 comprising the 714bp NruI - XhoI fragment of pRHP6, including part of the first exon, the intron, and the entire coding sequences of ICP47.
7. A vector comprising a virus including a vector element according to claim 5.
8. An vector according to claim 7 comprising a vector having the identifying characteristics of AdICP47-1.
9. A method for inhibiting cell recognition by cytotoxic T lymphocytes, comprising introducing into said cells an isolated protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and fragments of any of the foregoing that are able to inhibit said recognition.
10. A method for the treatment of herpesvirus infections, comprising introducing into infected cells a nucleotide sequence that is at least in part complementary to the mRNA sequence encoding a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and biologically active fragments of any of the foregoing, wherein



the complementary portion of said nucleotide sequence is of sufficient length to inhibit the translation of said mRNA and thereby inhibit the production of said protein.

11. A method for the treatment of herpesvirus infections, comprising introducing into infected cells an antibody specific for a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to ICP47 of HSV-1, and antigenic fragments of any of the foregoing.
12. A method for identifying drugs useful in treating herpesvirus infections, comprising:
 - Establishing a model cell system that expresses a protein selected from the group of ICP47 of HSV type 1, IE 12 of HSV type 2, proteins that are more than 40% homologous to the ICP47 protein of HSV-1, and fragments of any of the foregoing that exhibit the functional characteristics of ICP47;
 - Adding amounts of candidate compounds to samples of said model cells; and
 - Testing said samples for a trait different from that observed in samples to which no such compound has been added, said trait being selected from the group of suppressed synthesis of the ICP47 homologue, decreased MHC class I protein processing, and increased CTL lysis.
13. A drug identified by the method of claim 12.
14. A method for the prevention and treatment of

disorders the effects of which are at least in part caused by autoimmune reactions, the method comprising introducing into a patient's cells a biomolecule selected from the group of an isolated nucleotide sequence encoding ICP47 of HSV type 1, an isolated nucleotide sequence encoding IE 12 of HSV type 2, isolated nucleotide sequences encoding proteins that are more than 40% homologous to ICP47 of HSV-1, the protein ICP47, the protein IE 12, proteins that are more than 40% homologous to ICP47, and therapeutically effective fragments of any of the foregoing.

15. The method according to claim 14 wherein the disorder is tissue and organ transplant rejection.
16. The method according to claim 14 wherein the disorder is diabetes.
17. The method according to claim 14 wherein the disorder is multiple sclerosis.
18. The method according to claim 14 wherein the disorder is arthritis.
19. The method according to claim 14 wherein the disorder is ocular herpesvirus infection.

FIG. 1






FIG. 2A




FIG. 2B

FIG. 2C

FIG. 3






FIG. 4

A



B



4 / 8
SUBSTITUTE SHEET

FIG. 5








FIG. 6



6 / 8

SUBSTITUTE SHEET


FIG. 7



7 / 8
SUBSTITUTE SHEET

FIG. 8

Lysis by anti-HCMV CTL clone 16E6



8 / 8

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 94/00657

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/38 C12N7/01 C12N15/63 A61K48/00 C07K14/035
A61K38/16 A61K39/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF VIROLOGY, vol.54, no.2, May 1985 pages 414 - 421 PERSSON, R.H. ET AL. 'Cells that constitutively express the Herpes Simplex Virus immediate-early protein ICP4 allow efficient activation of viral delayed-early genes in trans' cited in the application see Figure 1; page 415, left-hand column, lines 43-49	5,6
Y	---	7,8
	--- --/--	

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not
considered to be of particular relevance"E" earlier document but published on or after the international
filing date"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)"O" document referring to an oral disclosure, use, exhibition or
other means"P" document published prior to the international filing date but
later than the priority date claimed"T" later document published after the international filing date
or priority date and not in conflict with the application but
cited to understand the principle or theory underlying the
invention"X" document of particular relevance; the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step when the document is taken alone"Y" document of particular relevance; the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 April 1995

Date of mailing of the international search report

18-05-1995

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Alt, G

Form PCT/ISA/210 (second sheet) (July 1992)

page 1 of 2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 94/00657

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>VIROLOGY, vol.195, no.1, July 1993, USA pages 62 - 70 BONNEAU, R.H. ET AL. 'Epitope specificity fo H-2Kb-restricted, HSV-1-, and HSV-2-cross-reactive cytotoxic T lymphocyte clones' see page 63, "Cells and Viruses" ---</p>	7,8
A	<p>JOURNAL OF VIROLOGY, vol.66, no.11, November 1992 pages 6264 - 6272 POSAVAD, C.M. AND ROSENTHAL, K.L. 'Herpes Simplex Virus-infected human fibroblasts are resistant to and inhibit cytotoxic T-lymphocyte activity' ---</p>	1
A	<p>J.GEN.VIROL., vol.67, 1986 pages 1035 - 1048 UMENE, K. 'Conversion of a fraction of the unique sequence to part of the inverted repeats of the S component of the Herpes Simplex Virus type 1 genome' cited in the application ---</p>	1
A	<p>VIROLOGY, vol.194, May 1993 pages 419 - 423 NISHIYAMA, Y. ET AL. 'The US9, 10, 11, and 12 genes of Herpes Simplex Virus type 1 are of no importance for its neurovirulence and latency in mice' ---</p>	1
P,X	<p>CELL,, vol.77, 20 May 1994 pages 525 - 535 YORK, I.A. ET AL. 'A cytosolic Herpes Simplex Virus protein inhibits antigen presentation to CD8+ T lymphocytes' ---</p>	1-19
P,X	<p>VIROLOGY, vol.200, no.1, April 1994, USA pages 236 - 245 BANKS, T.A. ET AL. 'Vaccination with the immediate-early protein ICP47 of Herpes Simplex Virus-type 1 (HSV-1) induces virus-specific lymphoproliferation, but fails to protect against lethat challenge' see page 242, right-hand column, third paragraph -----</p>	1-19

1

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

THIS PAGE BLANK (USPTO)